

# ORGOGOZO STROKE SCALE

Patient Name: \_\_\_\_\_

Rater Name: \_\_\_\_\_

Date: \_\_\_\_\_

Activity	Score
----------	-------

## CONSCIOUSNESS

- 0 = coma
- 5 = stupor
- 10 = drowsiness
- 15 = normal

\_\_\_\_\_

## VERBAL COMMUNICATION

- 0 = impossible
- 5 = difficult
- 10 = normal

\_\_\_\_\_

## EYES AND HEAD SHIFT

- 0 = forced
- 5 = gaze failure
- 10 = none

\_\_\_\_\_

## FACIAL MOVEMENTS

- 0 = paralysis
- 5 = normal

\_\_\_\_\_

## ARM RAISING

- 0 = impossible
- 5 = incomplete
- 10 = possible

\_\_\_\_\_

## HAND MOVEMENTS

- 0 = useless
- 5 = useful
- 10 = skilled
- 15 = normal

\_\_\_\_\_

## UPPER LIMB TONE

- 0 = increased or decreased
- 5 = normal

\_\_\_\_\_

## LEG RAISING

- 0 = impossible
- 5 = gravity
- 10 = resistance
- 15 = normal

\_\_\_\_\_

## FOOT DORSIFLEXION

- 0 = foot drop
- 5 = gravity
- 10 = resistance or normal

\_\_\_\_\_

## LOWER LIMB TONE

- 0 = increased or decreased
- 5 = normal

\_\_\_\_\_

**TOTAL (0-100):** \_\_\_\_\_

## References

---

Orgogozo JM, Capildeo R. "Development of a neurological score for clinical evaluation of infarctions in the Sylvian territory."  
*Presse Med* 1983;12(48):3039-44.